

UCSF

UC San Francisco Previously Published Works

Title

First Draft Genome Sequences of Neisseria sp. Strain 83E34 and Neisseria sp. Strain 74A18, Previously Identified as CDC Eugonic Fermenter 4b Species.

Permalink

<https://escholarship.org/uc/item/0xh5b6p1>

Journal

Genome announcements, 4(6)

ISSN

2169-8287

Authors

Greninger, Alexander L
Streithorst, Jessica
Chiu, Charles Y
et al.

Publication Date

2016-11-01

DOI

10.1128/genomea.01277-16

Peer reviewed

First Draft Genome Sequences of *Neisseria* sp. Strain 83E34 and *Neisseria* sp. Strain 74A18, Previously Identified as CDC Eugonic Fermenter 4b Species

Alexander L. Greninger, Jessica Streithorst,  Charles Y. Chiu, Steve Miller

Department of Laboratory Medicine, UCSF, San Francisco, California, USA

We report the first draft genome sequences of two isolates previously classified as CDC EF-4b species, *Neisseria* sp. 83E34 and *Neisseria* sp. 74A18. Both strains were isolated from patients with animal bites and likely constitute novel genomospecies with average nucleotide identities of <95% to other sequenced strains.

Received 20 September 2016 Accepted 22 September 2016 Published 10 November 2016

Citation Greninger AL, Streithorst J, Chiu CY, Miller S. 2016. First draft genome sequences of *Neisseria* sp. strain 83E34 and *Neisseria* sp. strain 74A18, previously identified as CDC eugonic fermenter 4b species. *Genome Announc* 4(6):e01277-16 doi:10.1128/genomeA.01277-16.

Copyright © 2016 Greninger et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Steve Miller, steve.miller@ucsf.edu.

CDC eugonic fermenter 4b (EF-4b) is a designation of Gram-negative bacteria that are members of animal oral flora and are typically isolated from animal bite wounds in humans (1). In 2006, this group was identified as *Neisseria zoodegmatis* based on 16S sequence and biochemical testing (1–3). Only five isolates from human infections have been reported since the taxonomical classification of *Neisseria zoodegmatis*, while >30 isolates have been collected from CDC EF-4b bacteria (1, 4, 5). CDC EF-4b has also been cultured from a case of infectious tenosynovitis due to a Siberian tiger bite (6). To date, the 16S sequence is the only available nucleotide sequence for *Neisseria zoodegmatis*.

We sequenced the first draft genomes of two bacterial isolates that had been identified as CDC group EF-4b from the University of California, San Francisco (UCSF) microbiology lab. Isolate 74A18 was isolated from a right-hand index finger culture of a patient with flexor tenosynovitis following a dog bite. Isolate 83E34 was isolated from a left-lateral-hand culture of a patient with cellulitis following a cat bite. Both patients also had *Pasteurella multocida* isolated concurrently from the wound cultures.

DNA was extracted using the Qiagen EZ1 DNA tissue kit. Nextera XT paired-end and Nextera mate-pair libraries were sequenced at 250/350 bp and 2 × 80 bp on an Illumina MiSeq, respectively, and on an Oxford Nanopore MinION sequencer. Illumina sequences were adapter and quality (Q20) trimmed using cutadapt or NxTrim, *de novo* assembled using SPAdes version 3.5, metagenomically screened for contaminating sequence with SURPI, and annotated via Prokka version 1.1 (7–12). A total of 6,878,276 (74A18) and 9,380,478 (83E34) paired-end reads and 4,052,930 (74A18) and 5,428,958 (83E34) mate-pair reads were recovered after trimming. *De novo* assembly yielded 40 contigs totaling 2,419,010 bp, with an N_{50} of 233,412 bp for *Neisseria* sp. 83E34, while *Neisseria* sp. 74A18 yielded 88 contigs totaling 2,572,932 bp, with an N_{50} of 53,927 bp.

BLASTN analysis of the 16S sequence from *Neisseria* sp. 74A18 showed 98.6% identity to *Neisseria shayegani* clone TM092 (accession no. KM462144) and 98.1% identity to *Neisseria zoodegmatis* strain N15a (accession no. JQ979306). BLASTN analysis of the

16S sequence from *Neisseria* sp. 83E34 demonstrated 99.3% identity to *Neisseria canis* oral taxon 137 (accession no. JN713302), 98.2% identity to *Neisseria shayegani* clone TM092 (accession no. KM462144), and 96.7% identity to *Neisseria zoodegmatis* strain N15a (accession no. JQ979306). However, alignment of *Neisseria* sp. 83E34 to *Neisseria canis* housekeeping genes revealed poor alignment (84.7% for *rpoB* [accession no. KM438030] and 88.4% for *cpn60* [accession no. KJ872773]), suggesting it may not be a strain of *Neisseria canis*. Pairwise whole-genome alignment by LASTZ revealed 89.3% identity between our two sequenced species and 93.9% identity between *Neisseria* sp. 83E34 and *Neisseria wadsworthii* 9715 (WGS AGAZ01), the closest whole genome available (13). These data are consistent with a high degree of genomic variability within related *Neisseria* spp. and isolates previously identified as CDC EF-4 members (14). Further sequencing of EF-4b species, including *Neisseria zoodegmatis*, will be needed to fully classify these species.

Accession number(s). These whole-genome shotgun projects have been deposited at DDBJ/ENA/GenBank under the accession numbers [LGYH000000000](https://www.ncbi.nlm.nih.gov/nuclseq/LGYH000000000/) (*Neisseria* sp. 83E34) and [LGZA000000000](https://www.ncbi.nlm.nih.gov/nuclseq/LGZA000000000/) (*Neisseria* sp. 74A18).

ACKNOWLEDGMENT

This work was supported by the University of California, San Francisco, Department of Laboratory Medicine.

REFERENCES

1. Vandamme P, Holmes B, Bercovier H, Coenye T. 2006. Classification of Centers for Disease Control Group eugonic fermenter (EF)-4a and EF-4b as *Neisseria animaloris* sp. nov. and *Neisseria zoodegmatis* sp. nov., respectively. *Int J Syst Evol Microbiol* 56:1801–1805. <http://dx.doi.org/10.1099/ijs.0.64142-0>.
2. Rossau R, Vandenbussche G, Thielemans S, Segers P, Grosch H, Gothe E, Mannheim W, De Ley J. 1989. Ribosomal ribonucleic acid cistron similarities and deoxyribonucleic acid homologies of *Neisseria*, *Kingella*, *Eikenella*, *Simonsiella*, *Alysiella*, and Centers for Disease Control groups EF-4 and M-5 in the emended family *Neisseriaceae*. *Int J Syst Bacteriol* 39:185–198. <http://dx.doi.org/10.1099/00207713-39-2-185>.
3. Barrett SJ, Sneath PHA. 1994. A numerical phenotypic taxonomic study

- of the genus *Neisseria*. Microbiology 140:2867–2891. <http://dx.doi.org/10.1099/00221287-140-10-2867>.
4. Heydecke A, Andersson B, Holmdahl T, Melhus A. 2013. Human wound infections caused by *Neisseria animaloris* and *Neisseria zoodegmatidis*, former CDC group EF-4a and EF-4b. Infect Ecol Epidemiol 3:20312. <http://dx.doi.org/10.3402/iee.v3i0.20312>.
 5. Ganière JP, Escande F, André-Fontaine G, Larrat M, Filloneau C. 1995. Characterization of group EF-4 bacteria from the oral cavity of dogs. Vet Microbiol 44:1–9. [http://dx.doi.org/10.1016/0378-1135\(94\)00110-1](http://dx.doi.org/10.1016/0378-1135(94)00110-1).
 6. Isotalo PA, Edgar D, Toye B. 2000. Polymicrobial tenosynovitis with *Pasteurella multocida* and other Gram negative bacilli after a Siberian tiger bite. J Clin Pathol 53:871–872. <http://dx.doi.org/10.1136/jcp.53.11.871>.
 7. Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. Bioinformatics 30:2068–2069. <http://dx.doi.org/10.1093/bioinformatics/btu153>.
 8. Naccache SN, Federman S, Veeraraghavan N, Zaharia M, Lee D, Samayoa E, Bouquet J, Greninger AL, Luk K-C, Enge B, Wadford DA, Messenger SL, Genrich GL, Pellegrino K, Grard G, Leroy E, Schneider BS, Fair JN, Martínez MA, Isa P, Crump JA, DeRisi JL, Sittler T, Hackett J, Miller S, Chiu CY. 2014. A cloud-compatible bioinformatics pipeline for ultrarapid pathogen identification from next-generation sequencing of clinical samples. Genome Res 24:1180–1192. <http://dx.doi.org/10.1101/gr.171934.113>.
 9. Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads. EMBnet J 17:10–12. <http://dx.doi.org/10.14806/ej.17.1.200>.
 10. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. J Comput Biol 19:455–477. <http://dx.doi.org/10.1089/cmb.2012.0021>.
 11. Greninger AL, Langelier C, Cunningham G, Keh C, Melgar M, Chiu CY, Miller S. 2015. Two rapidly growing mycobacterial species isolated from a brain abscess: first whole genome sequence of *Mycobacterium immunogenium* and *Mycobacterium llatzerense*. J Clin Microbiol 53:2374–2377. <http://dx.doi.org/10.1128/JCM.00402-15>.
 12. Greninger AL, Chorny I, Knowles S, Ng VL, Chaturvedi V. 2015. Draft genome sequences of four NDM-1-producing *Klebsiella pneumoniae* strains from a health care facility in northern California. Genome Announc 3(3):e00421-15. <http://dx.doi.org/10.1128/genomeA.00421-15>.
 13. Harris RS. 2007. Improved pairwise alignment of genomic DNA. Ph.D. thesis. Pennsylvania State University, Computer Science and Engineering, State College, PA.
 14. Roach DJ, Burton JN, Lee C, Stackhouse B, Butler-Wu SM, Cookson BT, Shendure J, Salipante SJ. 2015. A year of infection in the intensive care unit: prospective whole genome sequencing of bacterial clinical isolates reveals cryptic transmissions and novel microbiota. PLoS Genet 11:e1005413. <http://dx.doi.org/10.1371/journal.pgen.1005413>.